

## Featured Article

- Biotic invasion, niche stability, and the assembly of regional biotas in deep time: comparison between faunal provinces 359  
*Mark E. Patzkowsky and Steven M. Holland*

## Articles

- Quantitative analysis of the ecological dominance of benthic disaster taxa in the aftermath of the end-Permian mass extinction 380  
*Elizabeth Petsios and David J. Bottjer*
- Drilling predation increased in response to changing environments in the Caribbean Neogene 394  
*Jill S. Leonard-Pingel and Jeremy B. C. Jackson*
- Continental-scale biogeographic variation: provinces versus gradients in the Upper Ordovician of Laurentia 410  
*Chelsea E. Jenkins and Steven M. Holland*
- Axial growth gradients across the postprotaspid ontogeny of the Silurian trilobite *Aulacopleura koninckii* 426  
*Giuseppe Fusco, Paul S. Hong, and Nigel C. Hughes*
- The evolution of growth patterns in mammalian versus nonmammalian cynodonts 439  
*Rachel N. O'Meara and Robert J. Asber*
- A case study of extant and extinct Xenarthra cranium covariance structure: implications and applications to paleontology 465  
*Alex Hubbe, Diogo Melo, and Gabriel Marroig*
- Feeding ecology and habitat preferences of top predators from two Miocene carnivore-rich assemblages 489  
*M. Soledad Domingo, Laura Domingo, Juan Abella, Alberto Valenciano, Catherine Badgley, and Jorge Morales*
- Ecomorphological determinations in the absence of living analogues: the predatory behavior of the marsupial lion (*Thylacoleo carnifex*) as revealed by elbow joint morphology 508  
*Borja Figueirido, Alberto Martín-Serra, and Christine M. Janis*
- Body-size trends in *Peromyscus* (Rodentia: Cricetidae) on Vancouver Island, Canada, with comments on relictual gigantism 532  
*Martina L. Steffen*